


Spectroscopic Detection of Organophosphorus Agents


Dr. Roger M. Leblanc
Department of Chemistry
Coral Gables, FL

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 15 NOV 2004		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Spectroscopic Detection of Organophosphorus Agents				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Chemistry Coral Gables, FL				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM001849, 2004 Scientific Conference on Chemical and Biological Defense Research. Held in Hunt Valley, Maryland on 15-17 November 2004 . , The original document contains color images.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 26	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			


Insecticides and pesticides



Diethyl p-nitrophenyl phosphate
(Paraoxon)



Diethyl para-nitrophenol thiophosphate
(Parathion)




Di-isopropyl fluoro-phosphate
(DFP)


Nerve gases



Phosphonofluoridic acid, methyl-, 1,2,2-trimethylpropyl ester
(Soman)




Methylphosphonofluoridic acid 1-methyl-ethyl ester
(Sarin)



O-Ethyl S-(2-(diisopropylamino)ethyl)methylphosphonothioate
(VX)

Organophosphorus Hydrolase



- Isolated from *Pseudomonas diminuta*
- Metalloenzyme (zinc at active site)
- S_N2 mechanism of hydrolysis

Quantum Dot Characteristics

- Quantum dots
 - Size dependent emission
 - High quantum yield
 - Broad excitation spectra
 - Narrow emission band
 - Chemical versatility
- Core-shell quantum dots
 - Higher quantum yield
 - Chemical and photophysical stability
 - Low toxicity




1.9 2.4 4.1 5.2 nm



Excited with same
wavelength (365 nm)

Synthesis of Quantum Dots




Adam-Peng, Z. and Peng, X.
J. Am. Chem. Soc. **2001**, 123, 183-184

Chan, W. C. W. and Nie, S.
Science **1998**, 281, 2016-2018


Sensing device 1: CdSe composite films : paraoxon detection

- ✓ Layer-by-Layer technique (electrostatic interaction)





Constantine, C. A.; Gattás-Asfura, K. M.; Mello, S. V.; Crespo, G.; Rastogi, V.; Cheng, T.-C.; DeFrank, J. J. and Leblanc, R. M. *Langmuir* 2003, 19, 9863-9867

Optical properties of the quantum dots in aqueous solution :







Monitoring growth of the composite film on the quartz slide :

(1-7 = # layers of quantum dots)



Constantine, C. A.; Gattás-Asfura, K. M.; Mello, S. V.; Crespo, G.; Rastogi, V.; Cheng, T.-C.; DeFrank, J. J. and Leblanc, R. M. *Langmuir* 2003, 19, 9863-9867

Response of film upon exposure to paraoxon solution




Epifluorescence images of film

895μm × 713 μm



Before exposure
to paraoxon




After exposure
to paraoxon

Conclusion




- Quantum dots were successfully immobilized into composite films through utilization of the Layer-by-Layer technique and electrostatic interactions
- The convenient quantum dot-OPH composite film fabricated was highly selective and sensitive towards organophosphorus compounds (e.g. paraoxon)

Sensing device 2: Formation of OPH/QDs bioconjugate




Formation of OPH/QDs bioconjugation

Synthesis of CdSe(ZnS) QDs




PL spectra of OPH/QDs




Photoluminescence spectra of 10:1 molar ratio OPH/QDs bioconjugate in different concentrations of paraoxon. Samples were excited at 350 nm.

PL spectra of pure QDs




Photoluminescence spectra of pure QDs in different concentrations of paraoxon. Samples were excited at 350 nm.

Mechanism of PL quenching



Michaelis-Menten Equation



Steady state:

$$d[\text{ES}]/dt = k_{+1}(e_0 - [\text{ES}])[S] - k_{-1}[\text{ES}] - k_{+2}[\text{ES}] = 0 \quad ([S] = s_0 - [\text{ES}])$$



$$\longrightarrow [\text{ES}] = \frac{k_{+1} e_0 [S]}{k_{-1} + k_{+2} + k_{+1}[S]}$$

$$\longrightarrow [\text{ES}] = \frac{e_0 s_0}{(k_{-1} + k_{+2}) / k_{+1} + s_0} \quad (s_0 \gg e_0 > [\text{ES}])$$

$$\longrightarrow Q = k_q[\text{ES}] = \frac{k_q e_0 s_0}{(k_{-1} + k_{+2}) / k_{+1} + s_0}$$

Mechanism of PL quenching

Relative PL intensity percentage of (a) 10:1 (molar ratio) and (b) 100:1 (molar ratio) OPH/QDs bioconjugates as a function of paraoxon concentration.




F_0 : PL intensity at 573 nm in the absence of paraoxon
 F_n : PL intensity at 573 nm in the presence of paraoxon




Conclusion

- The OPH and CdSe(ZnS) QDs can form stable bioconjugate through electrostatic interaction
- CD spectra indicate a secondary structure change of OPH in the presence of paraoxon
- The intensity of photoluminescence of OPH/QDs bioconjugate was quenched in the presence of paraoxon
- The secondary structure change of OPH was responsible for the observed PL quenching
- Increasing the molar ratio of OPH over QDs will not substantially increase the sensitivity of OPH/QDs biosensor


Sensing device 3: OPH covalently bound to solid support




7-isothiocyanato-4-methylcoumarin (1)




Imaging ellipsometry




2-D image of the
silanized slide



3-D image of the
silanized slide




2-D image of
covalently attached OPH




3-D image of covalently
attached OPH

ATR-IR spectrum of the secondary structure of OPH





Frequency band position (cm ⁻¹)	Assignment
1693	anti-parallel β -sheet or pleated turn
1679	β -sheet
1657	α -helix
1647	amide I
1634	β -sheet
1619	β -sheet
1550	α -helix
1534	amide II

UV-Vis spectra for paraoxon analysis




Fluorescence spectra of OPH-coumarin *sensing system* in presence of paraoxon solution



Conclusion

- OPH was successfully bound to a silanized quartz substrate
- This bio-functional surface successfully detects the presence of paraoxon aqueous solution

Group Members



- Ph.D. graduate students
Celeste A. Constantine
Kerim M. Gattás-Asfura
Xiaojun Ji
Changqing Li
David Naistat
Jianmin Xu
Liang Zhao
Jiayin Zheng
- Undergraduate students
Gema Crespo
Philip DePaola
Andrea Escobar
Michael J. Snedaker
Daniel A. Thimann
Sanjeev Sirpal
- Senior Research Assoc.
Jhony Orbulescu

Acknowledgement

- U.S. Army Research Office
–(DAAD 19-03-1-0131)

